

ANTIMICROBIËLE DOSERINGEN BIJ KINDEREN				
Product	Route	Totale dagdosis (mg/kg)	Dosis-interval	Maximale dagdosis
amikacine	iv, im	15-20 mg/kg	1 x /d	1g
amoxicilline	po	50-100 mg/kg	3x /d	6g
amoxicilline/ clavulaanzuur	po	40-60 mg/kg	3x /d	4g
amoxicilline/ clavulaanzuur	iv	100-150 mg/kg	4 x /d	6-8g
amoxicilline	iv,im	100-200 mg/kg	4 x /d	8g
amoxicilline (hoge dosering)	iv,im	200-300 mg/kg	4 x /d	8g
cefazoline	iv,im	100-150 mg/kg	4 x /d	6g
cefepim	iv,im	150 mg/kg	3 x /d	6g
cefotaxim	iv,im	100-200 mg/kg	4 x /d	10g
ceftazidim	iv,im	150 mg/kg	3x /d	6g
ceftriaxon	iv,im	50-100 mg/kg	1-2 x /d	4g
cefuroxim	iv,im	100 mg/kg	4 x /d	6g
cefuroxim axetil	po	30-50 mg/kg	2dd	1,5g
ciprofloxacin	po	30 mg/kg	2x /d	1,5g
ciprofloxacin	iv	20 mg/kg	2 x /d	1,2g
clarithromycine	po	15 mg/kg	2 x /d	1g
clindamycine	po	20 mg/kg	4 x /d	1,8g
clindamycine	iv,im	25-40 mg/kg	4 x /d	2,4g
flucloxacilline	po	50-100 mg/kg	4 x /d	2g
flucloxacilline	iv,im	100-150 mg/kg	4 x /d	8g
linezolid	po,iv	30 mg/kg	3x /d	1,2g
meropenem	iv	60-120 mg/kg	3x /d	6g
metronidazol	po/iv	30 mg/kg	3-4 x /d	1,5g
penicilline G	iv,im	100.000-200.000 IE	4 x /d	20x106 IE g
penicilline G (hoge dosering)	iv,im	300.000 IE	6x /d	20x106 IE g
penicilline V	po	50.000 IE	4 x /d	4x106 IE g
piperacilline/ tazobactam	iv	200-300 mg/kg	4 x /d	16g
temocilline	iv,im	50 mg/kg	2 x /d	4g
vancomycine	iv	40-60 mg/kg	4 x /d	2g